

REPORT DOCUMENTATION PAGE			Form Approved OMB NO. 0704-0188	
Public Reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimates or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188,) Washington, DC 20503.				
1. AGENCY USE ONLY (Leave Blank)		2. REPORT DATE 10/1/04		3. REPORT TYPE AND DATES COVERED FINAL 01 Apr 02 – 31 Dec 03
4. TITLE AND SUBTITLE Electronic Wavefunction Imaging and Spectroscopy in Metallic and Magnetic Nanostructures by Millikelvin Scanning Tunneling Microscopy			5. FUNDING NUMBERS DAAD19-02-1-0043	
6. AUTHOR(S) J. C. Davis				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California, Berkeley, CA 94720, USA. Cornell University, Ithaca, NY 14850. USA.			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U. S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211			10. SPONSORING / MONITORING AGENCY REPORT NUMBER 43360.1-PH-QC	
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.				
12 a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			12 b. DISTRIBUTION CODE .	
13. ABSTRACT (Maximum 200 words) We developed a mK scanning tunneling microscope whereby we can interrogate the wavefunction of donor states at individual dopant atoms to determines if they are useful as qubits.				
14. SUBJECT TERMS			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OR REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION ON THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

Foreword

We use high resolution STM at millikelvin temperatures (mK-STM) for allow imaging, spectroscopy, and control of the quantum wavefunctions of the electrons at individual dopant atoms in superconductors and semiconductors.

Statement of Problem

To develop a millikelvin STM capable of operation at high-fields with atomic-resolution spectroscopic mapping and its application for wavefunction imaging studies of electronic states at individual dopant atoms in several types of semiconductors.

Summary of Achievements

1. Ultra Low Vibration Laboratory at Cornell

We have completed construction and testing of a new ultra low vibration (ULV) laboratory at Cornell that is specifically designed to facilitate these projects. It consists of an underground laboratory inside which are two nested acoustic isolation rooms. The inner room is supported on six vibration isolators. Inside the inner acoustic room is the cryostat itself which is made of ~3 tons of lead and houses another massive vibration isolation stage on three isolators. The dewar and refrigerator are suspended from this stage. ***This laboratory facility exceeds (see Fig. 1c below) the stringent vibration requirements set the proposed projects and is one of the few laboratories in the world that does so.***




Figure 1A. The schematic of overall ULV laboratory design at Cornell. This space is about 5 meters high and 6 m square




Figure 1B. The ULV lab+ cryostat performance. Measured vertical acceleration noise is suppressed into the $1 \text{ ngHz}^{-1/2}$ range for all frequencies above $\sim 10 \text{ Hz}$. The measurement under operating conditions (black) was made on board the refrigerator at $T=0.02 \text{ K}$ while the acoustic/RF enclosures were closed and the two sets of air springs activated. Very similar horizontal acceleration noise isolation is also detected.

2. Millikelvin STM Installation




Figure 2. The ULV lab+ STM cryostat inside the RF shield/acoustic shield room. Close-up fo Dilution refrigerator.





Fig.3 The STM head suspended below the mixing chamber at the center of the 9T magnet (not shown)

GaAs -Te Tests.

We used the millikelvin, high-field, scanning tunneling microscopy (STM) to study the fundamental physics of bound electronic states at individual dopant atoms in a semiconductor. Although much has been said and written about electronically addressing single quantum states at dopant atoms in semiconductors, no experiments have been carried out and almost nothing is known about the applied physics of this situation. Therefore our studies were initially directed towards direct detection and study of electronic bound-states at individual dopant Te atoms (in high magnetic fields at very low temperatures) as a test case. We could identify the location of the dopant atoms, measure their spectrum and locally map the wavefunction of the donor state with spectroscopic mapping. We chose GaAs because it cleaves very nicely on the $\langle 110 \rangle$ plane and has a direct band gap which is observable in tunneling spectroscopy.



GaAs (110) ideal / reconstructed surface

Fig. 4 Schematic of $\langle 110 \rangle$ cleave surface of GaAs




Fig. 5 Images of $\langle 110 \rangle$ surface of GaAs showing surface electronic defect states and the location of the Te donor atom.

When cleaved in cryogenic ultra high vacuum the surface has a small coverage of defect states which are possibly due to missing GaAs atoms from the cleave or perhaps by displacement of the atoms from their correct locations in the reconstruction. In addition to these phenomena, we can see the locations of the Te dopant atoms as faint circular regions of higher LDOS. They reside approximately 1.5 nm under the surface on average.

We demonstrated spectroscopic imaging of the wavefunction of the donor state. This is a *qua non* of direct manipulation of such states and the first time it has been achieved to our knowledge.

Publications

1. Relating atomic scale electronic phenomena to wave-like quasiparticle states in superconducting $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$, K. McElroy, R. W. Simmonds, J. E. Hoffman, D.-H. Lee, J. Orenstein, H. Eisaki, S. Uchida & J.C. Davis” **Nature** **422**, 520 (2003).
2. Incommensurate, dispersive, density of states modulations in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$, K. McElroy, J. E. Hoffman, D. -H. Lee, K. M. Lang, H. Eisaki, S. Uchida and J. C. Davis.” **Physica C** **388-389**, 225-226 (2003)
3. Vortex-induced quasi-particle ‘checkerboard’ in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$, J. E. Hoffman, E. W. Hudson, K. M. Lang, H. Eisaki, S. Uchida and J. C. Davis. **Physica C**, **388-389**, 703-704 (2003).
4. Numerical Studies of the Superfluid Shapiro Effect”. R. W. Simmonds, A. Marchenkov, J. C. Davis and R. E. Packard. **Physica B** **329**, 63(2003).

5. STM studies of individual Ti impurity atoms in Sr_2RuO_4 ”, Barker BI, Dutta SK, Lupien C, McEuen PL, Kikugawa N, Maeno Y, Davis JC **PHYSICA B-329**: 1334-1335 (2003)
6. Fourier Transform Scanning Tunneling Spectroscopy Studies of the Electronic Structure of Superconducting $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+d}$ ”, K. McElroy, J. E. Hoffman H. Eisaki S. Uchida & J.C. Davis, **AIP CONFERENCE PROCEEDINGS 696**, 1 (2003).
7. Fourier transform scanning tunneling spectroscopy: A new window on the electronic structure of $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+d}$ ” Davis JC. **ACTA. PHYS. POL. A 104** (3-4): 193-204 (2003).
8. Measurements of attenuation of third sound: Evidence of trapped vorticity in thick films of superfluid He-4 Hoffman, J A, Penanen K, Davis J.C. et al **J LOW TEMP PHYS 135**: 177-202 (2004)..
9. A ‘checkerboard’ electronic crystal state in in Lightly Hole-Doped $\text{Ca}_{2-x}\text{Na}_x\text{CuO}_2\text{Cl}_2$ ” T. Hanaguri, C. Lupien, Y. Kohsaka, D.-H. Lee, M. Azuma, M. Takano, H. Takagi, & J. C. Davis. **Nature 430**, 1001 (2004).

Personnel

Dr. Christian Lupien (moving to Cornell University to my new lab in June 2003).
Mr. Andy Schmidt.